



Detection of mcr-1 encoding plasmid-mediated colistin-resistant *Escherichia coli* isolates from human bloodstream infection and imported chicken meat, Denmark 2015

Hasman, H.; Hammerum, A. M.; Hansen, F.; Hendriksen, Rene S.; Olesen, B.; Agersø, Yvonne; Zankari, Ea; Leekitcharoenphon, Pimlapas; Stegger, M.; Kaas, Rolf Sommer

Total number of authors:
14

Published in:
Eurosurveillance (Online Edition)

Link to article, DOI:
[10.2807/1560-7917.es.2015.20.49.30085](https://doi.org/10.2807/1560-7917.es.2015.20.49.30085)

Publication date:
2015

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Hasman, H., Hammerum, A. M., Hansen, F., Hendriksen, R. S., Olesen, B., Agersø, Y., Zankari, E., Leekitcharoenphon, P., Stegger, M., Kaas, R. S., Cavaco, L., Hansen, D. S., Aarestrup, F. M., & Skov, R. L. (2015). Detection of mcr-1 encoding plasmid-mediated colistin-resistant *Escherichia coli* isolates from human bloodstream infection and imported chicken meat, Denmark 2015. *Eurosurveillance (Online Edition)*, 20(49), 1-5. <https://doi.org/10.2807/1560-7917.es.2015.20.49.30085>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Detection of *mcr-1* encoding plasmid-mediated colistin-resistant *Escherichia coli* isolates from human bloodstream infection and imported chicken meat, Denmark 2015

H Hasman¹, AM Hammerum¹, F Hansen¹, RS Hendriksen², B Olesen³, Y Agersø², E Zankari², P Leekitcharoenphon², M Stegger^{1,4}, RS Kaas², LM Cavaco², DS Hansen³, FM Aarestrup², RL Skov¹

1. Department of Microbiology and Infection Control, Statens Serum Institut, Copenhagen, Denmark

2. National Food Institute, Technical University of Denmark, Lyngby, Denmark

3. Department of Clinical Microbiology, Herlev and Gentofte Hospital, Copenhagen University Hospital, Herlev, Denmark

4. Pathogen Genomics Division, Translational Genomics Research Institute (TGen), Flagstaff, Arizona, USA

Correspondence: Henrik Hasman (henh@ssi.dk)

Citation style for this article:

Hasman H, Hammerum A, Hansen F, Hendriksen R, Olesen B, Agersø Y, Zankari E, Leekitcharoenphon P, Stegger M, Kaas R, Cavaco L, Hansen D, Aarestrup F, Skov R. Detection of *mcr-1* encoding plasmid-mediated colistin-resistant *Escherichia coli* isolates from human bloodstream infection and imported chicken meat, Denmark 2015. Euro Surveill. 2015;20(49):pii=30085. DOI: <http://dx.doi.org/10.2807/1560-7917.ES.2015.20.49.30085>

Article submitted on 04 December 2015 / accepted on 10 December 2015 / published on 10 December 2015

The plasmid-mediated colistin resistance gene, *mcr-1*, was detected in an *Escherichia coli* isolate from a Danish patient with bloodstream infection and in five *E. coli* isolates from imported chicken meat. One isolate from chicken meat belonged to the epidemic spreading sequence type ST131. In addition to *IncI2**, an *incX4* replicon was found to be linked to *mcr-1*. This report follows a recent detection of *mcr-1* in *E. coli* from animals, food and humans in China.

Very recently, in November 2015, Liu *et al.* reported the finding of a transferable plasmid-mediated colistin resistance gene, *mcr-1*, detected in *Escherichia coli* isolates from animals, food and patients in China. Moreover, they found *mcr-1* in *Klebsiella pneumoniae* isolates from patients [1]. Horizontal gene transfer represents a paradigm shift in colistin resistance, which until then only was found to be mediated by chromosomal mutations and thus spread by vertical transmission.

National surveillance of antimicrobial resistance in food animals, food and humans in Denmark using whole genome sequencing

Since 2012, the national surveillance of antimicrobial resistance in food animals, food and humans in Denmark (www.DANMAP.org) has used whole-genome sequence (WGS) analysis for detection of resistance genes and multilocus sequence typing (MLST) using the open-access bioinformatic web-tools ResFinder and MLST, respectively from www.genomicepidemiology.org for characterisation of extended spectrum beta-lactamase (ESBL)- and AmpC-producing *E. coli*

isolates [2-4]. The *mcr-1* sequence from China was added on 24 November 2015 to the ResFinder database as soon as it was available from The National Center for Biotechnology Information (NCBI).

Investigation of presence of *mcr-1* in *E. coli* isolates from food animals, food and human bloodstream infections

The updated version of ResFinder was used to analyse the WGS data from ESBL- and AmpC-producing *E. coli* isolates from food animals and food for the years 2012 to 2014, as well as ESBL- and AmpC-producing *E. coli* isolates from human bloodstream infections, and carbapenemase-producing organisms (CPOs) from humans, from January 2014 to beginning of November 2015 (Table 1), for the presence of *mcr-1*. Furthermore, fluoroquinolone resistance determinants were investigated by searching manually for mutations in the GyrA, ParC and ParE. [5].

The *mcr-1* gene was detected in one *E. coli* isolate from a human bloodstream infection from 2015 and in five *E. coli* isolates obtained from chicken meat of European origin imported to Denmark from 2012, 2013 and 2014 (Table 2). None of the CPOs were positive for *mcr-1* (Table 1).

The patient infected with the *mcr-1*-positive *E. coli* was an elderly man with prostate cancer and repeated urinary tract infections with ESBL-producing *E. coli* resulting in four positive urine samples over five month prior to the bloodstream infection, all resistant to third generation cephalosporins, gentamicin, sulfamethoxazole, trimethoprim and ciprofloxacin (these isolates were not

TABLE 1

Numbers of extended spectrum beta-lactamase and AmpC-producing *E. coli* isolates obtained and analysed by WGS from chicken meat, humans and carbapenemase-producing isolates from humans tested for *mcr-1* using ResFinder, Denmark, November 2015 (n=914)

Isolate origin	No. of isolates analysed by WGS	No. of sequences positive for <i>mcr-1</i>
ESBL- and AmpC-producing <i>E. coli</i> isolates from Danish chicken meat (2012–2014)	125	0
ESBL- and AmpC-producing <i>E. coli</i> isolates from imported chicken meat (2012–2014)	255	5
ESBL- and AmpC-producing <i>E. coli</i> isolates from human bloodstream infections (January 2014–beginning of November 2015)	417	1
Carbapenemase-producing isolates from humans (January 2014–beginning of November 2015)	117	0

ESBL: extended spectrum beta-lactamase; No: number; WGS: whole-genome sequence.

TABLE 2

Genotypic characterisation of *mcr-1*-positive *E. coli* isolates, Denmark, November 2015 (n=6)

Isolate name	Origin	Year of detection	MLST	Resistance genes detected by ResFinder besides <i>mcr-1</i>	Detection of chromosomal mutations encoding resistance to quinolones
0412016126	Chicken meat	2012	ST359	<i>aadA1</i> , <i>aadA5</i> , <i>aph(3')-Ic</i> , <i>bla</i> _{CMY-2} , <i>bla</i> _{TEM-1B} , <i>dfrA1</i> , <i>strA</i> , <i>strB</i> , <i>sul1</i> , <i>sul2</i> , <i>tet(B)</i>	GyrA (S83L, D87N) ParC (E62K)
0412044854	Chicken meat	2012	ST48	<i>aadA1</i> , <i>bla</i> _{CMY-2} , <i>bla</i> _{TEM-1B} , <i>dfrA1</i> , <i>mph(B)</i> , <i>strA</i> , <i>strB</i> , <i>sul1</i> , <i>sul2</i> , <i>tet(A)</i>	GyrA (S83L)
0412049521	Chicken meat	2012	ST131	<i>aadA1</i> , <i>bla</i> _{SHV-12} , <i>bla</i> _{CMY-2} , <i>strA</i> , <i>strB</i> , <i>sul1</i> , <i>dfrA1</i> , <i>tet(A)</i>	ND
0413040864	Chicken meat	2013	ST1112	<i>aadA1</i> , <i>aadA2</i> , <i>bla</i> _{SHV-12} , <i>cmlA1</i> , <i>sul3</i> , <i>tet(A)</i>	ND
14042624	Chicken meat	2014	ST2063	<i>aadA1</i> , <i>aadA2</i> , <i>bla</i> _{SHV-12} , <i>cmlA1</i> , <i>sul3</i>	ND
ESBL20150072	Human, bloodstream infection	2015	ST744	<i>aadA5</i> , <i>bla</i> _{CMY-2} , <i>bla</i> _{CTX-M-55} , <i>bla</i> _{TEM-1B} , <i>catA1</i> , <i>dfrA17</i> , <i>floR</i> , <i>fosA</i> , <i>mph(A)</i> , <i>rmtB</i> , <i>strA</i> , <i>strB</i> , <i>sul1</i> , <i>sul2</i> , <i>tet(A)</i>	GyrA (S83L, D87N) ParC (E62K)

MLST: multilocus sequence typing; ND: not detected.

investigated further). He had been treated empirically with piperacillin/tazobactam and subsequently meropenem after susceptibility testing of the bloodstream isolate, but not with colistin according to the available patient data.

Besides *mcr-1*, the human isolate from the Danish patient contained 15 different resistance genes including *bla*_{CTX-M-55} and *bla*_{CMY-2} conferring resistance to extended-spectrum beta-lactam antibiotics as well as two GyrA mutations (S83L, D87N) and a ParC mutation (E62K) leading to high-level fluoroquinolone resistance (Table 2). The human *mcr-1* positive *E. coli* isolate belonged to ST744, a rare sequence type in both humans and animals in Denmark. The patient had not been travelling abroad recently and the origin of the isolate is unknown.

The *bla*_{CMY-2} gene was detected in three of the five *mcr-1*-positive chicken meat isolates. In addition, three of

the chicken meat *E. coli* isolates carried *bla*_{SHV-12} conferring resistance to extended-spectrum beta-lactam antibiotics excluding cephamycins.

One of the *mcr-1* positive *E. coli* isolates from chicken meat belonged to ST131. The other chicken meat isolates belonged to sequence types not frequently found in Denmark (Table 2). The human MCR-1-producing *E. coli* isolate was only susceptible to piperacillin/tazobactam, carbapenems and tigecycline according to the European Committee on Antimicrobial Susceptibility Testing (EUCAST) breakpoints [6], whereas the chicken meat isolates were less resistant (Table 3).

WGS analysis using the web-tool PlasmidFinder [9] identified an IncI2 replicon present in the human isolate as well as in three of the chicken meat isolates, but this replicon was not detected in the remaining two chicken meat isolates. De novo assembly using CLCbio Genomics Workbench (v8.5.1; Qiagen, Aarhus,

TABLE 3

Antimicrobial susceptibility profiles of the five MCR-1-producing *E. coli* isolates from chicken meat and the MCR-1-producing *E. coli* isolate from human bloodstream infection, Denmark November 2015

Origin	Human		Chicken meat									
Isolate name	ESBL20150072		14042624		0413040864		0412049521		0412016126		0412044854	
Antimicrobial agent	MIC (mg/L)	S/R	MIC (mg/L)	S/R	MIC (mg/L)	S/R	MIC (mg/L)	S/R	MIC (mg/L)	S/R	MIC (mg/L)	S/R
Polymyxins												
Colistin	>4	R	>4	R	>4	R	>4	R	>4	R	>4	R
Polymyxin B ^a	4	R	>4	R	4	R	>4	R	>4	R	4	R
Beta-lactam/beta-lactam inhibitor combinations												
Ticarcillin/clavulanic acid	128/2	R	>128/2	R	≤16/2	S	64/2	R	64/2	R	≤16/2	S
Piperacillin/tazobactam	≤8/4	S	≤8/4	S	≤8/4	S	≤8/4	S	≤8/4	S	≤8/4	S
Cephalosporins												
Cefotaxime	>32	R	8	R	8	R	8	R	8	R	4	R
Ceftazidime	>16	R	>16	R	8	R	>16	R	16	R	8	R
Cefepime	>2	R	≤2	S	≤2	S	≤2	S	≤2	S	≤2	S
Monobactams												
Aztreonam	>16	R	>16	R	>16	R	>16	R	8	R	8	R
Carbapenems												
Ertapenem	≤0.25	S	≤0.25	S	≤0.25	S	≤0.25	S	≤0.25	S	≤0.25	S
Meropenem	≤1	S	≤1	S	≤1	S	≤1	S	≤1	S	≤1	S
Imipenem	≤1	S	≤1	S	≤1	S	≤1	S	≤1	S	≤1	S
Doripenem	≤0.125	S	≤0.125	S	≤0.125	S	≤0.125	S	≤0.125	S	≤0.125	S
Aminoglycosides												
Gentamicin	>8	R	≤1	S	2	S	≤1	S	≤1	S	≤1	S
Tobramycin	>8	R	≤1	S	2	S	≤1	S	2	S	≤1	S
Amikacin	>4	R	≤4	S	≤4	S	≤4	S	≤4	S	≤4	S
Fluoroquinolones												
Ciprofloxacin	>2	R	≤0.25	S	≤0.25	S	≤0.25	S	>2	R	≤0.25	S
Levofloxacin	8	R	≤1	S	≤1	S	≤1	S	>8	R	≤1	S
Tetracyclines												
Doxycycline ^b	8	I	≤2	S	16	R	8	I	16	R	8	I
Minocycline ^b	4	S	≤2	S	4	S	≤2	S	16	R	≤2	S
Glycylcyclines												
Tigecycline	≤0.25	S	≤0.25	S	≤0.25	S	≤0.25	S	≤0.25	S	≤0.25	S
Folate pathway inhibitors												
Trimethoprim/sulfamethoxazole	>4/76	R	≤0.5/9.5	S	≤0.5/9.5	S	>4/76	R	>4/76	R	>4/76	R

R/S according to the European Committee on Antimicrobial Susceptibility (EUCAST) clinical breakpoints [6].

I: intermediate; MIC: Minimal Inhibitory Concentration; R: resistant; S: sensitive.

^a Breakpoint according to Société Française de Microbiologie [7]

^b R/I/S according to The Clinical and Laboratory Standards Institute (CLSI) guidelines (M100-S25) [8].

Denmark) of the genomic data produced a direct link between the *mcr-1* gene and an IncX₄ replicon in one of the two isolates not containing IncI₂ replicons. An identical IncX₄ replicon was detected in four of the chicken meat isolates including both isolates lacking an IncI₂ replicon (but not in the human isolate).

Discussion and conclusion

Here we describe a MCR-1 producing *E. coli* isolate from a human infection coproducing both an ESBL (CTX-M-55) and an AmpC (CMY-2) cephalosporinase as well as five MCR-1 producing *E. coli* from chicken

meat coproducing either and ESBL (SHV-12) or an AmpC (CMY-2) cephalosporinase, or both. Human and animal CTX-M-55-producing isolates are commonly reported from Asia [10,11], but are relatively rarely seen in Denmark. CTX-M-55-producing *E. coli* isolates were detected in only 3% of the *E. coli* from bloodstream infections in 2014 [4]. CMY-2-producing *E. coli* isolates have commonly been detected from chicken meat both in Denmark and other countries [2-4,12,13], but *bla*_{CMY-2} has been relatively rare in the Danish human bloodstream *E. coli* isolates. Similarly, only two of the 245 human bloodstream *E. coli* isolates from 2014 were

SHV-12-producing [4]. Based on antibiogram data it seems plausible that the bloodstream infection is related to the repeated urinary tract infections, but this will need to be confirmed by additional WGS analysis. At this point in time, the origin of the human isolate is unresolved.

MLST analysis did not show any close clonal relationship between any of the six isolates. However, one of the chicken meat isolates belonged to ST131. This sequence type is commonly associated with human *E. coli* urinary tract and blood infection isolates worldwide, but are rare in animal *E. coli* isolates [4,14,15]. The fact that a ST131 MCR-1-producing *E. coli* isolate was found is of special concern, since ST131 isolates have spread epidemically during the last decade [14,15] and the ability of *mcr-1* to be acquired by this sequence type has been demonstrated here.

The *mcr-1* gene was initially reported to be located on an IncI2 plasmid without other known resistance markers [1]. Here only four of the isolates were found to contain an IncI2 replicon, suggesting that the *mcr-1* gene was either located on the chromosome or on a plasmid type belonging to another group. In support of the latter is the fact that *de novo* assembly of the genomic data from one of the isolates produced a continuous DNA fragment containing both an IncX4 and the *mcr-1* gene, strongly suggesting that the *mcr-1* gene is not restricted to the IncI2 plasmid group, but additional studies are needed to clarify this further.

In conclusion, this study is to our knowledge, the first proof of colistin-resistant *mcr-1* positive *E. coli* outside China. The human isolate was only susceptible to very few antimicrobial classes such as carbapenems. Should an isolate like this acquire carbapenem resistance, it would leave very few, if any, suitable treatment options. Finally, our findings underline the importance of continuous microbiological surveillance programs and not the least the benefit of employing comprehensive WGS-based surveillance of antimicrobial resistance, as it allows for rapid re-analysis of large datasets *in silico* and thus make early detection and risk assessment possible when new resistance genes emerge.

*Authors' correction

Upon request of the authors, Christina Aaby Svendsen's name was corrected in the Acknowledgement section on 14 December 2015. In addition, the sentence "In addition to IncN2, an incX4 replicon was found to be linked to *mcr-1*." was corrected to read "In addition to IncI2, an incX4 replicon was found to be linked to *mcr-1*." on 16 December 2015 upon request of the authors.

Acknowledgements*

The project was partly funded by the Danish Veterinary and Food Administration the centrally coordinated projects and partly by the Center for Genomic Epidemiology grant 09-067103/DSF from the Danish Council for Strategic Research

and supported by the Danish Ministry of Health as part of The Integrated Surveillance of ESBL/AmpC producing *E. coli* and Carbapenemase Producing Bacteria. We thank Karin Sindhøj Pedersen and Christina Aaby Svendsen for excellent technical assistance.

Conflict of interest

None declared.

Authors' contributions

HH and AMH collected the data and drafted the manuscript, HH, MS, PL, EZ and RK did the molecular analysis, FMA, FH, YA, RSH, LC, DSH, BO produced phenotypic data and participated in the coordination and concept of the manuscript, RLS coordinated and edited the manuscript.

References

1. Liu Y-Y, Wang Y, Walsh TR, Yi L-X, Zhang R, Spencer J, et al. Emergence of plasmid-mediated colistin resistance mechanism MCR-1 in animals and human beings in China: a microbiological and molecular biological study. *Lancet Infect Dis*. 2015;3099:1-8. .DOI: 10.3109/23744235.2015.1113556 PMID: 26603172
2. The Danish Integrated Antimicrobial Resistance Monitoring and Research Programme (DANMAP). DANMAP 2012-Use of antimicrobial agents and occurrence of antimicrobial resistance in bacteria from food animals, food and humans in Denmark. DANMAP: Copenhagen. Sep 2013. Available from: http://www.danmap.org/Downloads/~media/Projekt%20sites/Danmap/DANMAP%20reports/DANMAP%202012/Danmap_2012.ashx
3. The Danish Integrated Antimicrobial Resistance Monitoring and Research Programme (DANMAP). DANMAP 2013 - Use of antimicrobial agents and occurrence of antimicrobial resistance in bacteria from food animals, food and humans in Denmark. DANMAP: Copenhagen. Sep 2014. Available from: <http://www.food.dtu.dk/-/media/Institut/Foedevareinstitut/Publikationer/Pub-2014/Danmap-2013.ashx?la=da>
4. The Danish Integrated Antimicrobial Resistance Monitoring and Research Programme (DANMAP). DANMAP 2014 - Use of antimicrobial agents and occurrence of antimicrobial resistance in bacteria from food animals, food and humans in Denmark. DANMAP: Copenhagen. Sep 2015. Available from: <http://www.food.dtu.dk/-/media/Institut/Foedevareinstitut/Publikationer/Pub-2015/Rapport%20Danmap%202014.ashx?la=da>
5. Hopkins KL, Davies RH, Threlfall EJ. Mechanisms of quinolone resistance in *Escherichia coli* and *Salmonella*: recent developments. *Int J Antimicrob Agents*. 2005;25(5):358-73. .DOI: 10.1016/j.ijantimicag.2005.02.006 PMID: 15848289
6. European Committee on Antimicrobial Susceptibility Testing (EUCAST). [Internet]. [Accessed 1 Dec 2015]. Available from: <http://www.eucast.org>
7. Comité de l'Antibiogramme de la Société Française de Microbiologie Recommandations du CASFM, Communiqué 2014 (Edition May 2014). Available from: [http://www.sfm-microbiologie.org/UserFiles/files/casfm/CASFM_EUCAST_V1_o_2014\(1\).pdf](http://www.sfm-microbiologie.org/UserFiles/files/casfm/CASFM_EUCAST_V1_o_2014(1).pdf)
8. The Clinical and Laboratory Standards Institute (CLSI). [Internet]. [Accessed 1 Dec 2015]. Available from: <http://clsi.org/blog/2015/01/08/clsi-publishes-new-antimicrobial-susceptibility-testing-standards/>
9. Carattoli A, Zankari E, García-Fernández A, Voldby Larsen M, Lund O, Villa L, et al. In silico detection and typing of plasmids using PlasmidFinder and plasmid multilocus sequence typing. *Antimicrob Agents Chemother*. 2014;58(7):3895-903. .DOI: 10.1128/AAC.02412-14 PMID: 24777092
10. Tong P, Sun Y, Ji X, Du X, Guo X, Liu J, et al. Characterization of antimicrobial resistance and extended-spectrum β -lactamase genes in *Escherichia coli* isolated from chickens. *Foodborne Pathog Dis*. 2015;12(4):345-52. .DOI: 10.1089/fpd.2014.1857 PMID: 25785885
11. Xu G, An W, Wang H, Zhang X. Prevalence and characteristics of extended-spectrum β -lactamase genes in *Escherichia coli* isolated from piglets with post-weaning diarrhea in Heilongjiang province, China. *Front Microbiol*. 2015;6:1103. .DOI: 10.3389/fmicb.2015.01103 PMID: 26500640

12. Ben Sallem R, Ben Slama K, Sáenz Y, Rojo-Bezares B, Estepa V, Jouini A, et al. Prevalence and characterization of extended-spectrum beta-lactamase (ESBL)- and CMY-2-producing *Escherichia coli* isolates from healthy food-producing animals in Tunisia. *Foodborne Pathog Dis.* 2012;9(12):1137-42. .DOI: 10.1089/fpd.2012.1267 PMID: 23194332
13. National ESBL surveillance group, Leverstein-van Hall MA, Dierikx CM, Cohen Stuart J, Voets GM, van den Munckhof MP, van Essen-Zandbergen A, et al. . Dutch patients, retail chicken meat and poultry share the same ESBL genes, plasmids and strains. *Clin Microbiol Infect.* 2011;17(6):873-80. .DOI: 10.1111/j.1469-0691.2011.03497.x PMID: 21463397
14. Price LB, Johnson JR, Aziz M, Clabots C, Johnston B, Tchesnokova V, et al. The epidemic of extended-spectrum- β -lactamase-producing *Escherichia coli* ST131 is driven by a single highly pathogenic subclone, H30-Rx. *MBio.* 2013;4(6):e00377-13. .DOI: 10.1128/mBio.00377-13 PMID: 24345742
15. Peirano G, Pitout JD. Molecular epidemiology of *Escherichia coli* producing CTX-M beta-lactamases: the worldwide emergence of clone ST131 O25:H4. *Int J Antimicrob Agents.* 2010;35(4):316-21. .DOI: 10.1016/j.ijantimicag.2009.11.003 PMID: 20060273